


UPPER GASTROINTESTINAL TRACT ACID RELATED DISORDERS (UGITARD)



¹ Character, localization and duration of pain. Pain relation to meals and sleep. Use of alcohol, caffeine and tobacco. Stress or anxiety. ASA and NSAID'S use. Chronic renal failure, COPD, Cirrhosis. Renal transplant, hyperparathyroidism. Family History of PUD.

² Weight loss. Hematemesis or melena Dysphagia Anemia.

³ Education about irritants free diet (Caffeine, acids, condiments). Avoid alcohol, cigarettes, ASA, NSAID'S).

⁴ Some insurance companies requires Ba swallow/UGIS prior to referral.

⁵ Not useful to follow response to treatment.

⁶ Bismuth Subsalicylate 2 tabs. QID. +Metronidazole 500mg BID.+Amoxicillin 1 gm.BID.(Or Tetracilin 250mg QID for allergic PNC)+Ranitidine 150mg. BID. X two weeks.

⁷ Peptic Ulcer (4-8) weeks. Gastric Ulcer (4-12 weeks).

⁸ Ranitidine 150 BID, Metoclopramide 5mg AC/HS, Anti reflux measures X 4 weeks.

⁹ Prilosec 20mg/d or Protonix 40mg/day or Prevacid 30mg/day.

PPI-Se ordenará el más barato.

**Este Protocolo excluye pacientes pediátricos.